Brain Research, 545 (1991) 358–359
Elsevier

Author Index

Agnati, L.F., see Zoli, M., 199
Akiyama, K., see Daigen, A., 131
Aleman, D.O., see Carr, K.D., 283
Anderson, R.E., see Lyons, M.K., 339
Araki, T., see Hara, J., 87
Ashby Jr., C.R., see Edwards, E., 276

Bak, T.H., see Carr, K.D., 283
Belkin, M., see Schwartz, M., 334
Ben-Bassat, S., see Schwartz, M., 334
Beyreuther, K., see Kitamoto, T., 319
Biesold, D., see Webster, H.H., 292
Blake, M.J., see Mesco, E.R., 355
Bonhaus, D.W., Russell, R.D. and McNamara, J.O., Activation of substantia nigra pars reticulata neurons: role in the initiation and behavioral expression of kindled seizures, 41
Brückner, G., see Webster, H.H., 292
Burgess, G.M., see Dunn, P.M., 80

Cain, S.T., see Kasckow, J., 343
Carr, K.D., Aleman, D.O., Bak, T.H. and Simon, E.J., Effects of parabrachial opioid antagonism on stimulation-induced feeding, 283
Cohen, A., see Schwartz, M., 334
Coote, P.R., see Dunn, P.M., 80
Correa, F.M.A., Guilhaume, S.S. and Saaverda, J.M., Comparative quantification of rat brain and pituitary angiotensin-converting enzyme with autoradiographic and enzymatic methods, 215
Cserr, H.F., see Ichimura, T., 103
Cunningham, K.A., see Lee, M.A., 351

Daigen, A., Akiyama, K. and Otsuki, S., Long-lasting change in the membrane-associated protein kinase C activity in the hippocampal kindled rat, 131
Daikoku, S., see Hisano, S., 265
Date, I., Felten, S.Y. and Felten, D.L., The nigrostriatal dopaminergic system in MPTP-treated mice shows more prominent recovery by syngeneic adrenal medullary graft than by allogeneic or xenogeneic graft, 191
De Belleroche, J., see Wood, H., 183
De Moraes, V.Z., see Schmidt, S.L., 123
Dechesne, C.J., Hampson, D.R., Goping, G., Wheaton, K.D. and Wenthold, R.J., Identification and localization of a kainate binding protein in the frog inner ear by electron microscopy immunocytochemistry, 223
Depaulis, A., see Liu, Z., 1
Dodt, H.-U., Pawelzik, H. and Zieglgänsberger, W., Actions of noradrenaline on neocortical neurons in vitro, 307
Drucker-Colín, R., see Lavin, M.A., 164
Dunn, P.M., Coote, P.R., Wood, J.N., Burgess, G.M. and Rang, H.P., Bradykinin evoked depolarization of a novel neuroblastoma × DRG neurone hybrid cell line (ND7/23), 80
Dykes, R.W., see Webster, H.H., 292

Edwards, E., Ashby, Jr., C.R. and Wang, R.Y., The effect of typical and atypical antipsychotic drugs on the stimulation of phosphoinositide hydrolysis produced by the 5-HT_3 receptor agonist 2-methyl-serotonin, 276
Etoh, T., see Murakami, N., 347

Feller, D.J., Tso-Olivas, D.Y. and Savage, D.D., Hippocampal mossy fiber zinc deficit in mice genetically selected for ethanol withdrawal seizure susceptibility, 73
Felten, D.L., see Date, I., 191
Felten, S.Y., see Date, I., 191
Ferraguti, F., see Zoli, M., 199
Fitzgerald, R.S., see Shirahata, M., 297
Fraser, P.A., see Ichimura, T., 103
Fuxe, K., see Zoli, M., 199

Gold, M.S., see Morgan, M.M., 17
Goldsmith, C.H., see Witelson, S.F., 175
Goping, G., see Dechesne, C.J., 223
Govitrapong, P., see Phansuwan-Pujito, P., 49
Guilhaume, S.S., see Correa, F.M.A., 215
Gustafsson, J.-Å., see Zoli, M., 199

Hampson, D.R., see Dechesne, C.J., 223
Hapner, S.J., see Paden, C.M., 151
Hara, H., Onodera, H., Kato, H., Araki, T. and Kogure, K., Autoradiographic analysis of second messenger and neurotransmitter system receptors in the gerbil hippocampus following transient forebrain ischemia, 87
Hara, N., see Sawada, M., 248
Hilbich, C., see Kitamoto, T., 319
Hinch, D., see Van Reeth, O., 208
Hisano, S. and Daikoku, S., Existence of mutual synaptic relations between corticotropin-releasing factor-containing and somatostatin-containing neurons in the rat hypothalamus, 265
Huisman, E., see Kamphuis, W., 33

Ichimura, T., Fraser, P.A. and Cserr, H.F., Distribution of extracellular tracers in perivascular spaces of the rat brain, 103
Iwane, M., see Kiyota, Y., 322

Jennes, L., Dual projections of gonadotropin releasing hormone containing neurons to the interpeduncular nucleus and the vasculature in the female rat, 329
Joseph, J.A., see Mesco, E.R., 355

Kalivas, P.W., see Steketee, J.D., 287
Kamatchi, G.L. and Ticku, M.K., Tricyclic antidepressants inhibit Ca^{2+}-activated K^+-efflux in cultured spinal cord neurons, 59
Kamphuis, W., Huisman, E., Veerman, M.J. and Lopes da Silva, F.H., Development of changes in endogenous GABA release during kindling epileptogenesis in rat hippocampus, 33
Kasckow, J., Cain, S.T. and Nemeroff, C.A., Neurotensin effects on calcium/calmodulin-dependent protein phosphorylation in rat neostriatal slices, 343
Kato, H., see Hara, H., 87
Kellogg, C.K., see Primus, R.J., 257
Kitamoto, T., Muramoto, T., Hilbich, C., Beyreuther, K. and Tateishi, J., *N*-terminal sequence of prion protein is also integrated into kuru plaques in patients with Gerstmann-Sträussler syndrome, 319
Kiyota, Y., Takami, K., Iwane, M., Shino, A., Miyamoto, M., Tsukuda, R. and Nagaoka, A., Increase in basic fibroblast growth factor-like immunoreactivity in rat brain after forebrain ischemia, 322
Kogure, K., see Hara, H., 87
Kurata, M., Okuda, M., Muneyuki, M. and Ohnishi, S.T., [^{31}P]MRS study of the protective effects of prostaglandin oligomers on forebrain ischemia in rats, 315
Kuroda, H., see Murakami, N., 347
Kuschinsky, W., see Schröck, H., 234

Landgraf, R., Ramirez, A.D. and Ramirez, V.D., The positive feedback action of vasopressin on its own release from rat septal tissue in vitro is receptor-mediated, 137
Lavie, V., see Schwartz, M., 334
Lavín, M.A. and Drucker-Colín, R., Ontogeny of the electrophysiological activity of dopaminergic cells with special reference to the influence of adrenal medullary grafts on aging, 164
Lee, M.A., Paris, J.M. and Cunningham, K.A., Cocaine-induced behavioral sensitization is not associated with loss of GABA-immunoreactive neurons in the amygdala, 351
Lerchl, A., see Stokkan, K.-A., 66
Liebeskind, J.C., see Morgan, M.M., 17
Liu, X.-b, Zheng, Z.-h, Xi, M.-c. and Wu, C.-p., Distribution of synapses on fast and slow pyramidal tract neurons in the cat. An electron microscopic study, 239
Liu, Z., Vergnes, M., Depaulis, A. and Marescaux, C., Evidence for a critical role of GABAergic transmission within the thalamus in the genesis and control of absence seizures in the rat, 1
Lopes da Silva, F.H., see Kamphuis, W., 33
Lyons, M.K., Anderson, R.E. and Meyer, F.B., Corticotropin releasing factor antagonist reduces ischemic hippocampal neuronal injury, 339

Maeno, T., see Sawada, M., 248
Manhães, A.C., see Schmidt, S.L., 123
Marescaux, C., see Liu, Z., 1

Marunouchi, T., see Suzumura, A., 301
Mayhan, W.G., Responses of the basilar artery to products released by platelets during chronic hypertension, 97
McCann, U.D. and Ricaurte, G.A., Major metabolites of (±)3,4-methylenedioxyamphetamine (MDA) do not mediate its toxic effects on brain serotonin neurons, 279
McNamara, J.O., see Bonhaus, D.W., 41
Mesco, E.R., Joseph, J.A., Blake, M.J. and Roth, G.S., Loss of D_2 receptors during aging is partially due to decreased levels of mRNA, 355
Meyer, F.B., see Lyons, M.K., 339
Mikkelsen, J.D., see Phansuwan-Pujito, P., 49
Miyamoto, M., see Kiyota, Y., 322
Mody, I., see Otis, T.S., 142
Møller, M., see Phansuwan-Pujito, P., 49
Morgan, M.M., Gold, M.S., Liebeskind, J.C. and Stein, C., Periaqueductal gray stimulation produces a spinally mediated, opioid antinociception for the inflamed hindpaw of the rat, 17
Muneyuki, M., see Kurata, M., 315
Murakami, N., Takamure, M., Takahashi, K., Utunomiya, K., Kuroda, H. and Etoh, T., Long-term cultured neurons from rat suprachiasmatic nucleus retain the capacity for circadian oscillation of vasopressin release, 347
Muramoto, T., see Kitamoto, T., 319
Murray, T.F., see Steketee, J.D., 287

Nagaoka, A., see Kiyota, Y., 322
Narahashi, T., see Reuveny, E., 312
Nemeroff, C.A., see Kasckow, J., 343
Nonaka, K.O., see Stokkan, K.-A., 66

Ohnishi, S.T., see Kurata, M., 315
Okuda, M., see Kurata, M., 315
Onodera, H., see Hara, H., 87
Otis, T.S., Staley, K.J. and Mody, I., Perpetual inhibitory activity in mammalian brain slices generated by spontaneous GABA release, 142
Otsuki, S., see Daigen, A., 131

Paden, C.M. and Hapner, S.J., Monoclonal antibodies identify two novel proteins associated with vasopressin secretory granules of the rat neurohypophysis, 151
Paris, J.M., see Lee, M.A., 351
Pawelzik, H., see Dodt, H.-U., 307
Phansuwan-Pujito, P., Mikkelsen, J.D., Govitrapong, P. and Møller, M., A cholinergic innervation of the bovine pineal gland visualized by immunohistochemical detection of choline acetyltransferase-immunoreactive nerve fibers, 49
Pinard, E., see Roussell, S., 171
Primus, R.J. and Kellogg, C.K., Experience influences environmental modulation of function at the benzodiazepine (BZD)/GABA receptor chloride channel complex, 257

Ramirez, A.D., see Landgraf, R., 137
Ramirez, V.D., see Landgraf, R., 137
Rang, H.P., see Dunn, P.M., 80
Rasmusson, D.D., see Webster, H.H., 292
Reiter, R.J., see Stokkan, K.-A., 66
Reuveny, E. and Narahashi, T., Potent blocking action of lead on voltage-activated calcium channels in human neuroblastoma cells SH-SY5Y, 312
Ricaurte, G.A., see McCann, U.D., 279
Roberts, A., see Sillar, K.T., 24
Roth, G.S., see Mesco, E.R., 355
Roussel, S., Pinard, E. and Seylaz, J., Focal cerebral ischemia in chronic hypertension: no protection by (*R*)-phenylisopropyladenosine, 171
Russell, R.D., see Bonhaus, D.W., 41

Saaverda, J.M., see Correa, F.M.A., 215
Savage, D.D., see Feller, D.J., 73
Sawada, M., Hara, N. and Maeno, T., Ionic mechanism of the outward current induced by extracellular ejection of interleukin-1 onto identified neurons of *Aplysia*, 248
Schliebs, R., see Webster, H.H., 292
Schmidt, S.L., Manhães, A.C. and De Moraes, V.Z., The effects of total and partial callosal agenesis on the development of paw preference performance in the BALB/cCF mouse, 123
Schober, W., see Webster, H.H., 292
Schröck, H. and Kuschinsky, W., Effects of nicotine withdrawal on the local cerebral glucose utilization in conscious rats, 234
Schwartz, M., Solomon, A., Lavie, V., Ben-Bassat, S., Belkin, M. and Cohen, A., Tumor necrosis factor facilitates regeneration of injured central nervous system axons, 334
Sears, L.L. and Steinmetz, J.E., Dorsal accessory inferior olive activity diminishes during acquisition of the rabbit classically conditioned eyelid response, 114
Seylaz, J., see Roussel, S., 171
Shino, A., see Kiyota, Y., 322
Shiotsu, G., see Stevens, K.E., 8
Shirahata, M. and Fitzgerald, R.S., The presence of CO_2/HCO_3^- is essential for hypoxic chemotransduction in the in vivo perfused carotid body, 297
Sillar, K.T. and Roberts, A., Segregation of NMDA and non-NMDA receptors at separate synaptic contacts: evidence from spontaneous EPSPs in *Xenopus* embryo spinal neurons, 24
Simon, E.J., see Carr, K.D., 283
Solomon, A., see Schwartz, M., 334
Staley, K.J., see Otis, T.S., 142
Stein, C., see Morgan, M.M., 17
Stein, L., see Stevens, K.E., 8
Steinmetz, J.E., see Sears, L.L., 114
Steketee, J.D., Striplin, C.A., Murray, T.F. and Kalivas, P.W., Possible role for G-proteins in behavioral sensitization to cocaine, 287
Stevens, K.E., Shiotsu, G. and Stein, L., Hippocampal μ-receptors mediate opioid reinforcement in the CA3 region, 8
Stokkan, K.-A., Reiter, R.J., Nonaka, K.O., Lerchl, A., Yu, B.P. and Vaughan, M.K., Food restriction retards aging of the pineal gland, 66
Striplin, C.A., see Steketee, J.D., 287
Suzumura, A., Marunouchi, T. and Yamamoto, H., Morphological transformation of microglia in vitro, 301

Takahashi, K., see Murakami, N., 347
Takami, K., see Kiyota, Y., 322
Takamura, M., see Murakami, N., 347
Tateishi, J., see Kitamoto, T., 319
Tecco, J.M., see Van Reeth, O., 208
Ticku, M.K., see Kamatchi, G.L., 59
Toffano, G., see Zoli, M., 199
Tso-Olivas, D.Y., see Feller, D.J., 73
Tsukuda, R., see Kiyota, Y., 322
Turek, F.W., see Van Reeth, O., 208

Utunomiya, K., see Murakami, N., 347

Van Reeth, O., Hinch, D., Tecco, J.M. and Turek, F.W., The effects of short periods of immobilization on the hamster circadian clock, 208
Vaughan, M.K., see Stokkan, K.-A., 66
Veerman, M.J., see Kamphuis, W., 33
Vergnes, M., see Liu, Z., 1

Wang, R.Y., see Edwards, E., 276
Webster, H.H., Rasmusson, D.D, Dykes, R.W., Schliebs, R., Schober, W., Brückner, G. and Bieshold, D., Long-term enhancement of evoked potentials in raccoon somatosensory cortex following coactivation of the nucleus basalis of Meynert complex and cutaneous receptors, 292
Wenthold, R.J., see Dechesne, C.J., 223
Wheaton, K.D., see Dechesne, C.J., 223
Witelson, S.F. and Goldsmith, C.H., The relationship of hand preference to anatomy of the corpus callosum in men, 175
Wood, H. and De Belleroche, J., Induction of c-fos mRNA in cerebral cortex by excitotoxin stimulation of cortical inputs: involvement of *N*-methyl-D-aspartate receptors, 183
Wood, J.N., see Dunn, P.M., 80
Wu, C.-p., see Liu, X.-b., 239

Xi, M.-c., see Liu, X.-b, 239

Yamamoto, H., see Suzumura, A., 301
Yu, B.P., see Stokkan, K.-A., 66

Zheng, Z.-h., see Liu, X.-b., 239
Zieglgänsberger, W., see Dodt, H.-U., 307
Zoli, M., Ferraguti, F., Gustafsson, J.-A., Toffano, G., Fuxe, K. and Agnati, L.F., Selective reduction of glucocorticoid receptor immunoreactivity in the hippocampal formation and central amygdaloid nucleus of the aged rat, 199